



NQAC

Nestlé Quality Assurance Center  
Dublin

# Technical Datasheet

**Analysis Name:** Total Amino Acids by AccQ-Tag & UHPLC-UV

**Method Number:** LI-00.594

**Scope of Application:** Soy- and milk-based infant and adult formula and similar materials, infant cereals, dry and wet pet foods.

**Description:** Proteins are hydrolyzed with 6 M HCl for 24 h at 110°C. Phenol (0.1%) is added to prevent halogenation of tyrosine. 3,3'-Dithiodipropionic acid (DDP) is added to convert cystine and cysteine to S-2-carboxyethylthiocysteine (XCys) as described by Barkholt & Jensen (1989). The resulting derivative can be separated from other amino acids and used for quantification of cystine (Cys<sub>2</sub>).

After hydrolysis and neutralization, amino acids and converted cysteine (XCys) are derivatized with 6-aminoquinolyl-N-hydroxysuccinimidyl carbamate (AQC) using the AccQ•Tag Ultra Method (Waters Corporation, Milford, MA, USA). Derivatized amino acids are separated using reversed phase UHPLC with UV detection at 260 nm.

**Sample Weight Required:** 25 g

**Method Reference:** Barkholt V. and Jensen A. L. (1989). Amino acid analysis: Determination of cysteine plus half-cystine in proteins after hydrochloric acid hydrolysis with a disulphide compound as additive. *Analytical Biochemistry* 177, 318-322.

Waters Corporation (2007). UPLC amino acid analysis solution. System guide (71500129702/Revision B). Waters Corporation, Milford, MA, USA.

**Analytical Platform:** UPLC



NQAC

Nestlé Quality Assurance Center  
Dublin

**Special Information:** During acid hydrolysis, glutamine (Gln) and asparagine (Asn) are converted to glutamic acid (Glu) and aspartic acid (Asp), respectively. Thus, Glu values represent the combined values of Glu and Gln, and Asp values represent the combined values of Asp and Asn. Cys2 values represent the combined values of cysteine and cystine since both are converted to XCys by DDP.

Analyte Reported	Alias	Unit of Measure	Limit of Quantification	Reproducibility*
Histidine		g/100 g	0.02	
Taurine		mg/100 g	16	
Serine		g/100 g	0.01	
Arginine		g/100 g	0.02	
Glycine		g/100 g	0.01	
Aspartic Acid		g/100 g	0.02	
Glutamic Acid		g/100 g	0.02	
Threonine		g/100 g	0.02	
Alanine		g/100 g	0.01	
Proline		g/100 g	0.01	
Lysine		g/100 g	0.02	
Tyrosine		g/100 g	0.02	
Cystine		g/100 g	0.03	
Methionine		g/100 g	0.02	
Valine		g/100 g	0.02	
Isoleucine		g/100 g	0.02	
Leucine		g/100 g	0.02	
Phenylalanine		g/100 g	0.02	

\*Reproducibility: The relative difference between 2 independent single test results obtained using the same method, on identical test material by different operators at different days (for intermediate reproducibility test) should not be greater than:

- Infant formulas and infant cereals: 3 % to 16 % for all amino acids, except for Ser (up to 27%) and Cys2 (up to 39%)
- Dry and wet pet foods: 3% to 19% for all amino acids and 42% for Cys2

Which corresponds to the relative reproducibility limit, iR %, at 95% confidence level.